

Drug Monograph

[Drug Name](#) | [Mechanism of Action and Pharmacokinetics](#) | [Indications and Status](#) | [Adverse Effects](#) | [Dosing](#) | [Administration Guidelines](#) | [Special Precautions](#) | [Interactions](#) | [Recommended Clinical Monitoring](#) | [Supplementary Public Funding](#) | [References](#) | [Disclaimer](#)

A - Drug Name

palbociclib

COMMON TRADE NAME(S): Ibrance™

[back to top](#)

B - Mechanism of Action and Pharmacokinetics

Palbociclib is a selective, reversible small molecule inhibitor of cyclin-dependent kinases (CDK) 4 and 6. The drug inhibits cyclin D-CDK 4/6 complex activity, blocking cell cycle progression from G1 to S phase. Palbociclib in combination with an anti-estrogen agent inhibits cell proliferation and induction of cell senescence in estrogen receptor (ER) positive breast cancer models.

Absorption	Bioavailability	46% (mean)
	Peak plasma levels	6 to 12 hours (capsule); 4 to 12 hours (tablet)
	Time to reach steady state	8 days
	Effects with food	<p>Capsule: When administered with food, AUC and C_{max} increased (up to 21% and 38%, respectively) and exposure variability decreased. Palbociclib capsules should be taken with food.</p> <p>Tablet: When administered with food, AUC and C_{max} increased up to 22% and 26%, respectively; food had no significant impact on exposure variability. Palbociclib tablets may be taken with or without food.</p>

Distribution	PPB	85%
Metabolism	Palbociclib undergoes hepatic metabolism via oxidation and sulfonation, primarily by CYP3A and sulfotransferase (SULT2A1) enzymes; acylation and glucuronidation are minor metabolic pathways.	
	Inactive metabolites	Yes
Elimination	Half-life	29 hours (mean plasma)
	Feces	74%
	Urine	18%

[back to top](#)

C - Indications and Status

Health Canada Approvals:

- Breast cancer

(Includes conditional approvals)

Refer to the product monograph for a full list and details of approved indications.

[back to top](#)

D - Adverse Effects

Emetogenic Potential: Minimal – No routine prophylaxis; PRN recommended

The following adverse effects occurred in patients with breast cancer treated with palbociclib plus fulvestrant in a Phase III study, where the incidence was at least 2% greater than the placebo plus fulvestrant arm. The table also includes severe or life-threatening adverse effects from other sources or post-marketing.

ORGAN SITE	SIDE EFFECT* (%)	ONSET**
Cardiovascular	Venous thromboembolism (1%) (pulmonary embolism)	E

Dermatological	Alopecia (15%)	E
	Rash, pruritus (14%)	E
Gastrointestinal	Anorexia (13%)	E
	Constipation (17%)	E
	Diarrhea (19%)	E
	Mucositis (25%)	E
	Nausea, vomiting (29%)	E
General	Edema - limbs (8%)	E
	Fatigue (38%)	E
Hematological	Myelosuppression ± infection, bleeding (79%) (62% severe)	E
Nervous System	Dizziness (11%)	E
	Dysgeusia (6%)	E
	Headache (21%)	E
	Insomnia (11%)	E
Respiratory	Cough, dyspnea (13%)	E
	Pneumonitis (1%)	E

* "*Incidence*" may refer to an absolute value or the higher value from a reported range.
 "*Rare*" may refer to events with < 1% incidence, reported in post-marketing, phase 1 studies, isolated data or anecdotal reports.

** I = *immediate* (onset in hours to days) E = *early* (days to weeks)
 D = *delayed* (weeks to months) L = *late* (months to years)

The most common side effects for palbociclib include myelosuppression ± infection, bleeding, fatigue, nausea, vomiting, mucositis, headache, diarrhea, constipation, alopecia, rash, pruritus and anorexia.

Neutropenia was the most frequently reported adverse effect with a median onset of 15 days. Febrile neutropenia, including one fatal case, has been reported in 2% of patients across clinical trials.

Infections were reported more frequently. Patients should be warned of the increased risk of infection and promptly report any occurrences of fever to their health care team.

Severe or life-threatening **interstitial lung disease/pneumonitis** has been reported in combination with endocrine therapy in clinical trials and post-marketing (including fatal cases).

[back to top](#)

E - Dosing

Refer to protocol by which patient is being treated.

Pre- and perimenopausal women treated with palbociclib and an aromatase inhibitor or fulvestrant should also be treated with luteinizing hormone-releasing hormone (LHRH) agonists according to local clinical practice.

Adults:

Oral: 125 mg Daily on Days 1 to 21, every 28 days

in combination with an aromatase inhibitor (e.g., letrozole) or fulvestrant. Refer to regimen monographs for dosing details.

Dosage with Toxicity:

Dose Level	Palbociclib Dose (mg/day) for 3 out of 4 weeks
0	125
-1	100
-2	75
-3	If further dose reduction required, discontinue.

Toxicity	Grade	Palbociclib Dose
Hematologic	3	<p>Day 1: Hold and repeat CBC within 1 week. When recovered to Grade ≤ 2, re-start next cycle at same dose.</p> <p>Day 15 of 1st 2 cycles: Continue current dose to complete the cycle. Repeat CBC day 22.</p> <p>If Grade 4 on Day 22, see Grade 4 recommendation below.</p> <p>Consider dose reduction if > 1 week recovery or recurrent Grade 3 neutropenia in subsequent cycles.</p>

	3 with fever \geq 38.5°C and/or infection	Hold until recovery to Grade \leq 2. Restart at the next lower dose.
	4	Hold until recovery to Grade \leq 2. Restart at the next lower dose.
Symptoms of interstitial lung disease (ILD)/pneumonitis (treatment-related)	Any	Hold dose and investigate; discontinue if severe ILD confirmed.
Other non-hematologic	3 or 4 (if persisting despite medical treatment)	Hold until recovery to Grade \leq 1 or Grade \leq 2 (if not considered a safety risk). Restart at the next lower dose.

Dosage with Hepatic Impairment:

Mean fraction of unbound palbociclib in plasma increased with worsening hepatic function.

Hepatic Impairment	Starting Dose
Mild - Moderate (Child-Pugh class A and B)	No dosage adjustment needed.
Severe (Child-Pugh class C)	75 mg once daily (days 1 to 21; q28 days). Monitor for toxicity.

Dosage with Renal Impairment:

No adjustment is required for CrCl \geq 15 mL/min. There is no data available in patients requiring hemodialysis.

Dosage in the elderly:

No overall differences in efficacy were observed between patients aged 65 and older compared to younger patients. When combined with letrozole, patients \geq 65 were more likely to experience anemia.

Dosage based on gender:

Gender and body weight had no significant effect on drug exposure.

Dosage based on ethnicity:

No dose modification is required based on pharmacokinetic, safety and efficacy data across Asian and non-Asian populations.

Children:

The safety and efficacy of palbociclib has not been studied in children under 18 years.

[back to top](#)

F - Administration Guidelines

- Palbociclib **capsules** should be administered **with food**; palbociclib **tablets** may be given **with or without food**.
- Capsules or tablets should be swallowed whole and not chewed, crushed, opened, or split prior to administration.
- If a patient vomits or misses a dose, an extra dose should not be taken to make up for the vomited or missed dose. The next dose should be taken at the usual time.
- Grapefruit, pomegranate, starfruit, Seville oranges, their juices or products should be avoided during palbociclib treatment.
- **Capsules** should be stored at 20 to 25°C, with excursions permitted between 15 to 30°C. **Tablets** should be stored at 15 to 30°C in original packaging to protect from moisture.

[back to top](#)

G - Special Precautions

Contraindications:

- Patients who are hypersensitive to palbociclib or any ingredient in the formulation or component in the container.

Other Warnings/Precautions:

- As fatigue and dizziness have been reported, patients should exercise caution when driving or operating machinery.
- Capsules contain lactose; carefully consider use in patients with hereditary galactose intolerance, severe lactase deficiency or glucose-galactose malabsorption.

Other Drug Properties:

- Carcinogenicity: Unknown
An increased incidence of microglial cell tumors was observed male rats; the relevance to humans is unknown.

Pregnancy and Lactation:

- Genotoxicity: Probable
- Mutagenicity: No
- Fetotoxicity: Yes
Palbociclib is not recommended for use in pregnancy. Adequate contraception should be used by both sexes during treatment, and for at least **21 days** after the last dose (for females) and **97 days** after the last dose (for males).
- Excretion into breast milk: Unknown
Breastfeeding is not recommended.
- Fertility effects: Probable
Animal data suggest that palbociclib may affect male fertility. Sperm preservation should be considered prior to starting treatment in males.

[back to top](#)

H - Interactions

Palbociclib is a substrate and weak inhibitor of CYP3A and a moderate substrate of P-gp. Drug interactions are possible with strong CYP3A inducers and inhibitors. Palbociclib is not an inhibitor of CYP1A2, 2A6, 2B6, 2C9 and 2D6 and not an inducer of CYP1A2, 2B6, 2C8 and 3A4 in vitro.

The drug has a low potential to inhibit drug transporters P-gp, BCRP, OAT1, OAT3, OCT2, OATP1B1 and OATP1B3. Palbociclib is not a substrate of OATP1B1 and OATP1B3 (in vitro)

studies).

There are no drug interactions with letrozole or goserelin.

Palbociclib solubility is pH dependent. Drug interactions with proton pump inhibitors are minimized when palbociclib capsules are given with food; food also reduces drug exposure variability of palbociclib capsules.

AGENT	EFFECT	MECHANISM	MANAGEMENT
CYP3A inhibitors (i.e. ketoconazole, clarithromycin, ritonavir, fruit or juice from grapefruit, Seville oranges or starfruit)	↑ palbociclib concentration and/or toxicity	↓ metabolism of palbociclib	Avoid strong CYP3A inhibitors.
Strong and moderate CYP3A inducers (i.e. phenytoin, rifampin, dexamethasone, carbamazepine, phenobarbital, St. John's Wort, etc)	↓ palbociclib concentration and/or efficacy	↑ metabolism of palbociclib	Avoid CYP3A inducers. If use of moderate inducer cannot be avoided, no dose adjustment to palbociclib is needed.
Antacids and PPIs (e.g. rabeprazole)	↓ palbociclib exposure when PPI and palbociclib capsules given in fasted conditions; minimal effect when palbociclib capsules given with food	Palbociclib has pH-dependent solubility; reduced palbociclib capsule solubility with increasing pH.	To minimize interaction with antacids and PPIs, palbociclib capsules should be given with food.
CYP3A substrates (e.g. cyclosporine, pimeozide, tacrolimus, triazolo-benzodiazepines, dihydropyridine calcium-channel blockers, certain HMG-CoA reductase inhibitors)	↑ substrate concentration and/or toxicity	Palbociclib is a weak inhibitor of CYP3A.	Consider reducing the dose of CYP3A substrates with narrow therapeutic indices (e.g. cyclosporine)

[back to top](#)

I - Recommended Clinical Monitoring

Treating physicians may decide to monitor more or less frequently for individual patients but should always consider recommendations from the product monograph.

Recommended Clinical Monitoring

Monitor Type	Monitor Frequency
CBC	Baseline and before each cycle, on day 15 of the first 2 cycles, one week after Grade 3 neutropenia, and as clinically indicated. If neutropenia Grade 2 or less in the first 6 cycles, may monitor every 3rd cycle thereafter
Liver function tests	Baseline and as clinically indicated
Renal function tests	Baseline and as clinically indicated
Clinical toxicity assessment for infection, bleeding, thromboembolism, pneumonitis, rash, headache, mucositis, fatigue and GI effects	At each visit

Grade toxicity using the current [NCI-CTCAE \(Common Terminology Criteria for Adverse Events\) version](#)

[back to top](#)

J - Supplementary Public Funding

Exceptional Access Program ([EAP Website](#))

- palbociclib - For the treatment of patients with estrogen receptor (ER)-positive, human epidermal growth factor receptor 2 (HER 2)-negative, unresectable locally advanced or metastatic breast cancer in combination with an aromatase inhibitor or fulvestrant, according to clinical criteria

[back to top](#)

K - References

Finn RS, Crown JP, Lang I, et al. The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study. *Lancet Oncol.* 2015 Jan;16(1):25-35.

Product Monograph: Ibrance (palbociclib). Pfizer Canada Inc. January 24, 2020.

April 2021 Updated indication and dosing description

[back to top](#)

L - Disclaimer

Refer to the [New Drug Funding Program](#) or [Ontario Public Drug Programs](#) websites for the most up-to-date public funding information.

The information set out in the drug monographs, regimen monographs, appendices and symptom management information (for health professionals) contained in the Drug Formulary (the "Formulary") is intended for healthcare providers and is to be used for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or adverse effects of a particular drug, nor should it be construed to indicate that use of a particular drug is safe, appropriate or effective for a given condition. The information in the Formulary is not intended to constitute or be a substitute for medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.

The format and content of the drug monographs, regimen monographs, appendices and symptom management information contained in the Formulary will change as they are reviewed and revised on a periodic basis. The date of last revision will be visible on each page of the monograph and regimen. Since standards of usage are constantly evolving, it is advised that the Formulary not be used as the sole source of information. It is strongly recommended that original references or product monograph be consulted prior to using a chemotherapy regimen for the first time.

Some Formulary documents, such as the medication information sheets, regimen information sheets and symptom management information (for patients), are intended for patients. Patients should always consult with their healthcare provider if they have questions regarding any information set out in the Formulary documents.

While care has been taken in the preparation of the information contained in the Formulary, such information is provided on an "as-is" basis, without any representation, warranty, or condition, whether express, or implied, statutory or otherwise, as to the information's quality, accuracy, currency, completeness, or reliability.

CCO and the Formulary's content providers shall have no liability, whether direct, indirect, consequential, contingent, special, or incidental, related to or arising from the information in the Formulary or its use thereof, whether based on breach of contract or tort (including negligence), and even if advised of the possibility thereof. Anyone using the information in the Formulary does so at his or her own risk, and by using such information, agrees to indemnify CCO and its content providers from any and all liability, loss, damages, costs and expenses (including legal fees and expenses) arising from such person's use of the information in the Formulary.

[back to top](#)